

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Guinea Pig IgG (H+L)-Alk. Phos., MinX none, ALP, Polyclonal , AP DNA-SEC-182911

| Article Name | Goat IgG anti-Guinea Pig IgG (H+L)-Alk. Phos., MinX none, ALP, Polyclonal , AP |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-182911 |
| Supplier Catalog Number | SEC-182911 |
| Alternative Catalog Number | DNA-SEC-182911 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Guinea pig |
| Immunogen | Guinea Pig IgG whole molecule |
| Conjugation | Alk. Phos. |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Anti-Guinea Pig IgG Alkaline Phosphatase Antibody generated in goat detects guinea pig IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to virus |

| Clonality | Polyclonal |
|--------------------|--|
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Goat Serum, Guinea Pig IgG and Guinea Pig Serum. |
| Form | Liquid (sterile filtered) |
| Formula | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,1% NaN3 |
| Target | Guinea Pig |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,000 |
| Application Notes | Anti-Guinea Pig IgG Alk Phos conjugate is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency. |