

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit Anti-Golden Syrian Hamster IgG F(ab)2 Antibody Peroxidase Conjugated - 607-4304, HRP, Polyclonal DNA-SEC-182952

Article Name	Rabbit Anti-Golden Syrian Hamster IgG F(ab)2 Antibody Peroxidase Conjugated - 607-4304, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-182952
Supplier Catalog Number	DNA-SEC-182952
Alternative Catalog Number	DNA-SEC-182952
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Golden Hamster
Immunogen	Hamster IgG F(ab)2 fragment
Conjugation	HRP
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Golden Syrian Hamster IgG F(ab)2 Peroxidase Antibody generated in rabbit is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 mol...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Hamster IgG, Hamster IgG F(ab)2 and Hamster Serum. No reaction was observed against Hamster IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Golden Syrian Hamster
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Hamster IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenzthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:7,000 to 1:30,000 of the reconstitution concentration is suggested for this product.