



Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Human IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh, Polyclonal DNA-SEC-183000

Article Name	Goat IgG anti-Human IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh, Polyclonal
Biozol Catalog Number	DNA-SEC-183000
Supplier Catalog Number	SEC-183000
Alternative Catalog Number	DNA-SEC-183000
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine, Gallus, Goat, Guinea pig, Hamster (all), Equine, Mouse, Rabbit, Rat, Sheep
Product Description	Anti-Human IgG (H&L) Biotin generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains - two identical heavy chains and two identic

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Secondary antibody conjugate was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against antibiotin, anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat and Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Fluorochrome Protein Value: 10-20, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Biotin Secondary antibody reagents are ideal for western blotting, Immunohistochemistry and ELISA as well as other antibody detection methods.