

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Human Lambda light chain-HRPO, MinX none DNA-SEC-183027

Article Name	Goat IgG anti-Human Lambda light chain-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-183027
Supplier Catalog Number	SEC-183027
Alternative Catalog Number	DNA-SEC-183027
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Anti-Human lambda chain was produced by repeated immunization with human lambda light chain in goat.
Conjugation	HRPO
Format	IgG
Target Specificity	Lambda (light chain)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	The anti-Human lambda (lambda chain) Antibody detects the lambda chain subunit. Immunoglobulins are heterotetramers composed of 2 immunoglobulin heavy and 2 immunoglobulin light chains. The immunoglobulin light chain is the small polypeptide subunit ...
Clonality	Polyclonal

Concentration	1.0 mg/ml
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. Specificity was confirmed by ELISA minimal cross reactivity against other human heavy or light chain isotypes.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	This secondary peroxidase conjugated antibody anti-Human lambda chain is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. Human lambda chain Antibody peroxidase conjugated has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:16,000 to 1:64,000 is suggested for this product.