

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Anti-Human IgG (H&L) Antibody Biotin Conjugated - 609-1602, Polyclonal DNA-SEC-183040

Article Name	Goat Anti-Human IgG (H&L) Antibody Biotin Conjugated - 609-1602, Polyclonal
Biozol Catalog Number	DNA-SEC-183040
Supplier Catalog Number	DNA-SEC-183040
Alternative Catalog Number	DNA-SEC-183040
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Human IgG (H&L) Biotin generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains - two identical heavy chains and two identic...

Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Human IgG, and Human Serum.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Fluorochrome Protein Value: 10-20, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Human IgG (H&L) Biotin Conjugate has been tested by ELISA and is assayed against 1.0 ug of Human IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:100,000 to 1:400,000 of the reconstitution concentration is suggested for this product.