

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit IgG anti-Human IgG (F(ab)2)-Alk. Phos., MinX Ms, ALP, Polyclonal , AP DNA-SEC-183084

Article Name	Rabbit IgG anti-Human IgG (F(ab)2)-Alk. Phos., MinX Ms, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-183084
Supplier Catalog Number	SEC-183084
Alternative Catalog Number	DNA-SEC-183084
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG F(ab)2 fragment
Conjugation	Alk. Phos.
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	Mouse
Product Description	Anti-Human IgG F(ab)2 Alkaline Phosphatase Antibody generated in rabbit recognizes the dimeric Fab portion of the human IgG molecule. Human IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme p

Clonality	Polyclonal
Concentration	1.4 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase (calf intestine), anti-Rabbit Serum, Human IgG, Human IgG F(ab')2 and Human Serum. No reaction was observed against Human IgG F(c) or Mouse Serum Proteins.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	This product has been assayed against 1.0 $\mu$ g of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,500 to 1:11,000 of the stated concentration is suggested for this product.