

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Mouse IgG (Fc)-HRPO, MinX none DNA-SEC-183171

Article Name	Goat IgG anti-Mouse IgG (Fc)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-183171
Supplier Catalog Number	SEC-183171
Alternative Catalog Number	DNA-SEC-183171
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Mouse
Immunogen	Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat.
Conjugation	HRPO
Format	IgG
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Mouse IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of mouse IgG and oft...
Clonality	Polyclonal

Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab') <sub>2</sub> fragment.
Form	Lyophilized
Formula	20 mM K <sub>3</sub> PO <sub>4</sub> , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Anti-Mouse IgG F(c) fragment is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.