

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit Anti-Mouse IgG (H&L) Antibody Peroxidase Conjugated - 610-4302, HRP, Polyclonal DNA-SEC-183266

Article Name	Rabbit Anti-Mouse IgG (H&L) Antibody Peroxidase Conjugated - 610-4302, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183266
Supplier Catalog Number	DNA-SEC-183266
Alternative Catalog Number	DNA-SEC-183266
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA, WB
Species Reactivity	Mouse
Immunogen	Mouse IgG whole molecule
Conjugation	HRP
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Secondary antibodies bind to the primary antibody to assist in detection, sorting and purification of target antigens. To enable detection, the secondary antibody must have specificity for the antibody species and isotype of the primary antibody bein...

Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	HRP Secondary Antibody Conjugate was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Mouse IgG and Mouse Serum.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:50,000 - 1:300,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:10,000 - 1:30,000
Application Notes	Anti-Mouse secondary antibody conjugated to horseradish peroxidase (HRP) generated in rabbit detects specifically Mouse IgG whole molecule. Anti-Mouse IgG peroxidase antibody has been tested by ELISA and western blot and is suitable for ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation, Immunohistochemistry as well as other HRP antibody based assays. Specific conditions for reactivity and signal detection should be optimized by the end user.