

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Mouse IgG (F(ab)2)-Biotin, MinX none DNA-SEC-183312

| | |
|----------------------------|---|
| Article Name | Rabbit IgG anti-Mouse IgG (F(ab)2)-Biotin, MinX none |
| Biozol Catalog Number | DNA-SEC-183312 |
| Supplier Catalog Number | SEC-183312 |
| Alternative Catalog Number | DNA-SEC-183312 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Mouse |
| Immunogen | Mouse IgG F(ab)2 fragment |
| Conjugation | Biotin |
| Format | IgG |
| Target Specificity | IgG (F(ab')2) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Anti-Mouse IgG F(ab)2 Biotin antibody generated in rabbit recognizes the dimeric Fab portion of the mouse IgG molecule. Mouse IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under co... |
| Clonality | Polyclonal |

| | |
|--------------------|--|
| Concentration | 2.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Mouse IgG, Mouse IgG F(ab') ₂ and Mouse Serum. No reaction was observed against Mouse IgG F(c). |
| Form | Lyophilized |
| Formula | 10 mM NaPO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Thimerosal |
| Target | Mouse |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | This product has been assayed against 1.0 µg of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. |