

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Donkey IgG anti-Mouse IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh DNA-SEC-183334

Article Name	Donkey IgG anti-Mouse IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh
Biozol Catalog Number	DNA-SEC-183334
Supplier Catalog Number	SEC-183334
Alternative Catalog Number	DNA-SEC-183334
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Mouse
Immunogen	Anti-Mouse IgG whole molecule was produced by repeated immunization with Mouse IgG whole molecule in donkey.
Conjugation	HRPO
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Rabbit,Rat,Sheep

Product Description	Anti-Mouse IgG whole molecule antibody generated in donkey detects specifically Mouse IgG whole molecule . This secondary antibody anti-Mouse is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immu...
Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Mouse IgG (H&L) Antibody Peroxidase Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:80,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:20,000
Application Notes	Suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other peroxidase antibody-based enzymatic assays requiring lot-to-lot consistency.