

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Rabbit IgG (F(ab)2)-HRPO, MinX Hu DNA-SEC-183371

|                            |   |
|----------------------------|---|
| Article Name               | Goat IgG anti-Rabbit IgG (F(ab)2)-HRPO, MinX Hu   |
| Biozol Catalog Number      | DNA-SEC-183371  |
| Supplier Catalog Number    | SEC-183371  |
| Alternative Catalog Number | DNA-SEC-183371  |
| Manufacturer               | dianova   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | ELISA,IHC,WB  |
| Species Reactivity         | Rabbit  |
| Immunogen                  | Anti-Rabbit IgG F(ab)2 fragment was produced by repeated immunization with Rabbit IgG F(ab)2 fragment in goat.  |
| Conjugation                | HRPO  |
| Format                     | IgG   |
| Target Specificity         | IgG (F(ab')2)   |
| Cross-Adsorption (MinX)    | Human   |
| Product Description        | Anti-Rabbit IgG F(ab)2 fragment antibody generated in goat detects specifically Rabbit IgG F(ab)2 fragment. This secondary antibody anti-Rabbit is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, im... |
| Clonality                  | Polyclonal  |

|                    |  |
|--------------------|--|
| Concentration      | 2.0 mg/mL  |
| Isotype            | Ig   |
| Buffer             | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
| Purity             | Anti-RABBIT IgG F(ab')2 (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab')2 and Rabbit Serum. No reaction was observed against Rabbit IgG F(c) or Human Serum Proteins. |
| Form               | Lyophilized  |
| Formula            | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target             | Rabbit   |
| Antibody Type      | Secondary Antibody   |
| Application Dilute | ELISA Dilution: 1:130,000, Immunohistochemistry Dilution: 1:1,000 - 1:10,000, Western Blot Dilution: 1:2,000 - 1:20,000  |
| Application Notes  | Anti-Rabbit IgG F(ab')2 fragment is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.  |