

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Rat IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rb,Sh DNA-SEC-183449

Article Name	Goat IgG anti-Rat IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rb,Sh
Biozol Catalog Number	DNA-SEC-183449
Supplier Catalog Number	SEC-183449
Alternative Catalog Number	DNA-SEC-183449
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rat
Immunogen	Rat IgG whole molecule
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Sheep
Product Description	Anti-Rat IgG (H&L) generated in goat detects rat Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the c...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4, 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.