

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX none DNA-SEC-183496

| Article Name | Rabbit IgG anti-Rat IgG (H+L)-FITC, MinX none |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-183496 |
| Supplier Catalog Number | SEC-183496 |
| Alternative Catalog Number | DNA-SEC-183496 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | FLISA,FACS,IF |
| Species Reactivity | Rat |
| Immunogen | Rat IgG whole molecule |
| Conjugation | FITC |
| Format | IgG |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Anti-Rat IgG (H&L) fluorescein Antibody generated in rabbit detects reactivity to Rat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruse |
| Clonality | Polyclonal |

| Concentration | 2.0 mg/mL |
|--------------------|--|
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | Anti-Rat IgG (H&L) fluorescein Antibody generated in rabbit detects reactivity to Rat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsinization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both the Heavy and Light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. This Anti-Rat IgG antibody is conjugated to Fluorescein. |
| Form | Lyophilized |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Rat |
| Antibody Type | Secondary Antibody |
| Application Dilute | FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 3.1, IF Microscopy Dilution: 1:1,000 - 1:5,000 |
| Application Notes | Secondary antibody reagents are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. |