

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Donkey IgG anti-Sheep IgG (H+L)-Biotin, MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt DNA-SEC-183572

Article Name	Donkey IgG anti-Sheep IgG (H+L)-Biotin, MinX Ck,Gp,Hm,Ho,Hu,Ms,Rb,Rt
Biozol Catalog Number	DNA-SEC-183572
Supplier Catalog Number	SEC-183572
Alternative Catalog Number	DNA-SEC-183572
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Sheep
Immunogen	Sheep IgG whole molecule
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Gallus,Guinea pig,Hamster (all),Equine,Human,Mouse,Rabbit,Rat
Product Description	Anti-Sheep IgG (H&L) generated in donkey detects sheep Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Donkey Serum, Sheep IgG and Sheep Serum. No reaction was observed against Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4, 150 mM NaCl, pH 7.2, lyophilisate, 0.01% NaN3
Target	Sheep
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Sheep IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 is suggested for this product.