

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Hamster (all) IgG (H+L)-Alk. Phos., MinX none, ALP, Polyclonal , AP DNA-SEC-183709

Article Name	Rabbit F(ab)2 anti-Hamster (all) IgG (H+L)-Alk. Phos., MinX none, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-183709
Supplier Catalog Number	SEC-183709
Alternative Catalog Number	DNA-SEC-183709
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Hamster (all)
Immunogen	Golden Syrian Hamster IgG whole molecule
Conjugation	Alk. Phos.
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Golden Syrian Hamster IgG Alkaline Phosphatase Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies fo...

Clonality	Polyclonal
Concentration	0.6 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Rabbit Serum, Golden Syrian Hamster IgG and Golden Syrian Hamster Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c). Limited reactivity will occur against Armenian Hamster IgG.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Golden Syrian Hamster
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:3,000 - 1:12,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 µg of Hamster IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.