

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat F(ab)2 Anti-Human IgG IgA IgM (H&L) Antibody Peroxidase Conjugated (Min X Mouse Serum Proteins) - 709-103-130, HRP, Polyclonal DNA-SEC-183720

Article Name	Goat F(ab)2 Anti-Human IgG IgA IgM (H&L) Antibody Peroxidase Conjugated (Min X Mouse Serum Proteins) - 709-103-130, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183720
Supplier Catalog Number	DNA-SEC-183720
Alternative Catalog Number	DNA-SEC-183720
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG, IgA, and IgM whole molecules
Conjugation	HRP
Format	F(ab')2
Target Specificity	IgG+IgM+IgA (H+L)
Cross-Adsorption (MinX)	Mouse
Product Description	F(ab)2 Anti-Human IgG IgA IgM (H&L) Peroxidase Antibody generated in goat detects human (heavy and light chain) immunoglobulin G, A, and M. Immunoglobulin G binds to antigens and can neutralize or opsonize targets, and are predominantly involved in t...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c) or Mouse Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.
Form	Lyophilized
Formula	20 mM K3PO4, 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	F(ab)2 Anti-Human IgG IgA IgM (H&L) Peroxidase Antibody has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.