

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 Anti-Human IgG (H&L) Antibody Peroxidase Conjugated (Min X Bv Hs Ms & Rt Serum Proteins) - 709-1316, HRP, Polyclonal DNA-SEC-183741

Article Name	Goat F(ab)2 Anti-Human IgG (H&L) Antibody Peroxidase Conjugated (Min X Bv Hs Ms & Rt Serum Proteins) - 709-1316, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183741
Supplier Catalog Number	DNA-SEC-183741
Alternative Catalog Number	DNA-SEC-183741
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	HRP
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Equine,Mouse,Rat
Product Description	F(ab)2 Anti-Human IgG (H&L) Peroxidase Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum i...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Horse, Mouse and Rat Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4, 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:50,000 - 1:125,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	F(ab)2 Anti-Human IgG (H&L) Peroxidase Antibody has been tested by ELISA and is assayed against 1.0 µg of Human IgG in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:20,000 of the reconstitution concentration is suggested for this product.