

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Human IgM (μ)-Alk. Phos., MinX none, ALP, Polyclonal , AP DNA-SEC-183751

Article Name	Goat F(ab)2 anti-Human IgM (μ)-Alk. Phos., MinX none, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-183751
Supplier Catalog Number	SEC-183751
Alternative Catalog Number	DNA-SEC-183751
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgM Fc5μ fragment
Conjugation	Alk. Phos.
Format	F(ab')2
Target Specificity	IgM (μ)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Anti-Human IgM Fc5μ Alkaline Phosphatase Antibody generated in goat detects specifically the Fc5μ portion of the human IgM heavy chain. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure t...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Human IgM and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Human IgG or Human IgA. Specificity was confirmed by ELISA at less than 1% cross reactivity against other human heavy or light chain isotypes.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN ₃
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:7,500 - 1:30,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. F(ab') ₂ Anti-Human IgM Fc5μ antibody has been assayed against 1.0 μg of Human IgM in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.