

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Donkey F(ab)2 anti-Human IgG (H+L)-unconj., MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh DNA-SEC-183762

Article Name	Donkey F(ab)2 anti-Human IgG (H+L)-unconj., MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh
Biozol Catalog Number	DNA-SEC-183762
Supplier Catalog Number	SEC-183762
Alternative Catalog Number	DNA-SEC-183762
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	Unconjugated
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Mouse,Rabbit,Rat,Sheep
Product Description	F(ab)2 Anti-Human IgG (H&L) Antibody generated in donkey detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglob...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Human IgG, and Human Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c), or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Mouse, Rabbit, Rat, and Sheep Serum Proteins.
Form	Liquid (sterile filtered)
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000, Immunohistochemistry Dilution: 1:1,000-1:5,000, Western Blot Dilution: 1:2,000-1:10,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1×10^6 cells in flow cytometry is approximately 1.0 μ g of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.