

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-FITC, MinX Hu, Polyclonal DNA-SEC-183772

Article Name	Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-FITC, MinX Hu, Polyclonal
Biozol Catalog Number	DNA-SEC-183772
Supplier Catalog Number	SEC-183772
Alternative Catalog Number	DNA-SEC-183772
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FACS,IF
Species Reactivity	Mouse
Immunogen	Mouse IgG IgA and IgM whole molecule
Conjugation	FITC
Format	F(ab')2
Target Specificity	IgG+IgM+IgA (H+L)
Cross-Adsorption (MinX)	Human
Product Description	F(ab)2 Anti-Mouse IgG IgA IgM (H&L) Fluorescein Antibody generated in goat detects reactivity to Mouse IgG, Mouse IgA, and Mouse IgM subclasses and chains. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing ...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-Pepsin or Human Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	Flow Cytometry Dilution: User Optimized, IF Microscopy Dilution: 1:500 - 1:2,500
Application Notes	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.