

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-HRPO, MinX Hu DNA-SEC-183774

Article Name	Goat F(ab)2 anti-Mouse IgG+IgM+IgA (H+L)-HRPO, MinX Hu
Biozol Catalog Number	DNA-SEC-183774
Supplier Catalog Number	SEC-183774
Alternative Catalog Number	DNA-SEC-183774
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Mouse
Immunogen	Anti-Mouse IgG IgA and IgM (H&L) was produced by repeated immunization with mouse IgG IgA and IgM molecule in goat.
Conjugation	HRPO
Format	F(ab')2
Target Specificity	IgG+IgM+IgA (H+L)
Cross-Adsorption (MinX)	Human
Product Description	F(ab)2 Anti-Mouse IgG IgA IgM (H&L) Peroxidase Antibody generated in goat detects reactivity to Mouse IgG, Mouse IgA, and Mouse IgM subclasses and chains. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-pepsin, or Human Serum Proteins. This reagent is suitable for the detection of all mouse subclasses, isotypes and chain combinations.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:80,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Antibodies Anti-Mouse IgG IgA IgM peroxidase conjugated are suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.