

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG (H+L)-FITC, MinX Bo,Ho,Hu,Rb,Rt,Sh DNA-SEC-183788

Article Name	Goat F(ab)2 anti-Mouse IgG (H+L)-FITC, MinX Bo,Ho,Hu,Rb,Rt,Sh
Biozol Catalog Number	DNA-SEC-183788
Supplier Catalog Number	SEC-183788
Alternative Catalog Number	DNA-SEC-183788
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FLISA,FACS,IF
Species Reactivity	Mouse
Immunogen	Mouse IgG whole molecule
Conjugation	FITC
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Equine,Human,Rabbit,Rat,Sheep
Product Description	F(ab)2 Anti-Mouse IgG (H&L) Fluorescein Antibody generated in goat was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Horse, Human, Rabbit, Rat or Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 2.8, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.