

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 Anti-Mouse IgG Fc Antibody Peroxidase Conjugated (Min X Bv Hs & Hu Serum Proteins) - 710-1322, HRP, Polyclonal DNA-SEC-183791

Article Name	Goat F(ab)2 Anti-Mouse IgG Fc Antibody Peroxidase Conjugated (Min X Bv Hs & Hu Serum Proteins) - 710-1322, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183791
Supplier Catalog Number	DNA-SEC-183791
Alternative Catalog Number	DNA-SEC-183791
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA
Species Reactivity	Mouse
Immunogen	Mouse IgG F(c) fragment
Conjugation	HRP
Format	F(ab')2
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	Bovine,Equine,Human
Product Description	F(ab)2 Anti-Mouse IgG F(c) Peroxidase Antibody was generated in goat and detects specifically Mouse IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration mu...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	F(ab)2 Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab) or Bovine, Horse, and Human Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:250 - 1:1000, Western Blot Dilution: 1:500 - 1:2,000
Application Notes	F(ab)2 Anti-Mouse IgG F(c) Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.