

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Mouse IgG (Fc)-HRPO, MinX Bo,Ho,Hu, Polyclonal DNA-SEC-183791

| | |
|----------------------------|---|
| Article Name | Goat F(ab)2 anti-Mouse IgG (Fc)-HRPO, MinX Bo,Ho,Hu, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183791 |
| Supplier Catalog Number | SEC-183791 |
| Alternative Catalog Number | DNA-SEC-183791 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Mouse |
| Immunogen | Mouse IgG F(c) fragment |
| Conjugation | HRPO |
| Format | F(ab')2 |
| Target Specificity | IgG (Fc) |
| Cross-Adsorption (MinX) | Bovine,Equine,Human |
| Product Description | F(ab)2 Anti-Mouse IgG F(c) Peroxidase Antibody was generated in goat and detects specifically Mouse IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration mu... |
| Clonality | Polyclonal |

| | |
|--------------------|---|
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | F(ab') ₂ Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(ab) or Bovine, Horse, and Human Serum Proteins. |
| Form | Lyophilized |
| Formula | 20 mM K ₃ PO ₄ , 150 mM NaCl, pH 7.2, lyophilisate, 0.01% Gentamicin |
| Target | Mouse |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:100,000, Immunohistochemistry Dilution: 1:250 - 1:1000, Western Blot Dilution: 1:500 - 1:2,000 |
| Application Notes | F(ab') ₂ Anti-Mouse IgG F(c) Antibody is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10 ⁶ cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |