

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 Anti-Mouse IgG F(ab)2 Antibody Alkaline Phosphatase Conjugated (Min X Bv Hs Hu Rb Rt & Sh Serum Proteins) - 710-1531, AP, Polyclonal DNA-SEC-183797

Article Name	Goat F(ab)2 Anti-Mouse IgG F(ab)2 Antibody Alkaline Phosphatase Conjugated (Min X Bv Hs Hu Rb Rt & Sh Serum Proteins) - 710-1531, AP, Polyclonal
Biozol Catalog Number	DNA-SEC-183797
Supplier Catalog Number	DNA-SEC-183797
Alternative Catalog Number	DNA-SEC-183797
Manufacturer	danova
Host	Goat
Category	Antikörper
Application	ELISA
Species Reactivity	Mouse
Immunogen	Mouse IgG F(ab)2 fragment
Conjugation	AP
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	Bovine,Equine,Human,Rabbit,Rat,Sheep

Product Description	F(ab)2 Anti-Mouse IgG F(ab)2 Alkaline Phosphatase Antibody generated in goat detects Mouse F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesi...
Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Mouse IgG, Mouse IgG F(ab)2 and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(c) or Bovine, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,09% NaN3
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	F(ab)2 Anti-Mouse IgG F(ab)2 Alkaline Phosphatase Antibody has been tested by ELISA and is assayed against 1.0 µg of Mouse IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:25,000 of the reconstitution concentration is suggested for this product. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.