

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Mouse IgG (H+L)-RPE, MinX Hu, Polyclonal DNA-SEC-183813

Article Name	Rabbit F(ab)2 anti-Mouse IgG (H+L)-RPE, MinX Hu, Polyclonal
Biozol Catalog Number	DNA-SEC-183813
Supplier Catalog Number	SEC-183813
Alternative Catalog Number	DNA-SEC-183813
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	FACS,IF
Species Reactivity	Mouse
Immunogen	Anti-Mouse IgG was produced by repeated immunization with Mouse IgG whole molecule in rabbit.
Conjugation	RPE
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Human
Product Description	F(ab)2 Mouse IgG (H&L) Antibody Phycoerythrin Conjugated Pre-Adsorbed was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies ...
Clonality	Polyclonal

Concentration	0.5 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Rabbit Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Human Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	Flow Cytometry Dilution: 1:100 - 1:250, IF Microscopy Dilution: 1:100 - 1:250
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1×10^6 cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.