

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Donkey F(ab)2 anti-Mouse IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh, Polyclonal DNA-SEC-183818

Article Name	Donkey F(ab)2 anti-Mouse IgG (H+L)-Biotin, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Rb,Rt,Sh, Polyclonal
Biozol Catalog Number	DNA-SEC-183818
Supplier Catalog Number	SEC-183818
Alternative Catalog Number	DNA-SEC-183818
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Mouse
Immunogen	Mouse IgG whole molecule
Conjugation	Biotin
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Rabbit,Rat,Sheep
Product Description	F(ab)2 Anti-Mouse IgG (H&L) Biotin Antibody generated in donkey was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for us...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Donkey Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Mouse
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:25,000 to 1:100,000 is suggested for this product.