

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 Anti-Rabbit IgG F(ab)2 Antibody (Min X Bv Hs Hu Ms Rt & Sh Serum Proteins) - 711-1124, Unconjugated, Polyclonal DNA-SEC-183828

Article Name	Goat F(ab)2 Anti-Rabbit IgG F(ab)2 Antibody (Min X Bv Hs Hu Ms Rt & Sh Serum Proteins) - 711-1124, Unconjugated, Polyclonal
Biozol Catalog Number	DNA-SEC-183828
Supplier Catalog Number	DNA-SEC-183828
Alternative Catalog Number	DNA-SEC-183828
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA, WB
Species Reactivity	Rabbit
Immunogen	Rabbit IgG F(ab)2 fragment
Conjugation	Unconjugated
Format	F(ab')2
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	Bovine,Equine,Human,Mouse,Rat,Sheep
Product Description	F(ab)2 Anti-Rabbit IgG F(ab)2 Antibody generated in goat detects Rabbit F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG, Rabbit IgG F(ab)2 and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rabbit IgG F(c) or Bovine, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
Form	Liquid (sterile filtered)
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:3,000, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	F(ab)2 Anti-Rabbit IgG F(ab)2 Antibody has been tested by ELISA and western blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.