

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat F(ab)2 anti-Rabbit IgG (H+L)-RPE, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh DNA-SEC-183841

Article Name	Goat F(ab)2 anti-Rabbit IgG (H+L)-RPE, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh
Biozol Catalog Number	DNA-SEC-183841
Supplier Catalog Number	SEC-183841
Alternative Catalog Number	DNA-SEC-183841
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FACS,IF
Species Reactivity	Rabbit
Immunogen	Rabbit IgG whole molecule
Conjugation	RPE
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Gallus,Goat,Guinea pig,Hamster (all),Equine,Human,Mouse,Rat,Sheep
Product Description	F(ab)2 Anti-Rabbit IgG (H&L) Antibody generated in goat detects immunoglobulin g from Rabbit, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglob...

Clonality	Polyclonal
Concentration	0.5 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Goat Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat and Sheep Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	Flow Cytometry Dilution: 1:100 - 1:250, IF Microscopy Dilution: 1:100 - 1:250
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody conjugate. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.