

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat F(ab)2 anti-Rat IgG (H+L)-unconj., MinX none DNA-SEC-183848

Article Name	Goat F(ab)2 anti-Rat IgG (H+L)-unconj., MinX none
Biozol Catalog Number	DNA-SEC-183848
Supplier Catalog Number	SEC-183848
Alternative Catalog Number	DNA-SEC-183848
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rat
Immunogen	Rat IgG whole molecule
Conjugation	Unconjugated
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper...
Clonality	Polyclonal

Concentration	1.03 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin and anti-Goat IgG F(c).
Form	Liquid (sterile filtered)
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.