

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (H+L)-unconj., MinX Bo, Ho, Hu, Ms, Rb, Sh DNA-SEC-183857

| | |
|----------------------------|---|
| Article Name | Goat F(ab)2 anti-Rat IgG (H+L)-unconj., MinX Bo, Ho, Hu, Ms, Rb, Sh |
| Biozol Catalog Number | DNA-SEC-183857 |
| Supplier Catalog Number | SEC-183857 |
| Alternative Catalog Number | DNA-SEC-183857 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA, IHC, WB |
| Species Reactivity | Rat |
| Immunogen | Rat IgG whole molecule |
| Conjugation | Unconjugated |
| Format | F(ab')2 |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | Bovine, Equine, Human, Mouse, Rabbit, Sheep |
| Product Description | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Clonality | Polyclonal |

| | |
|--------------------|--|
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Horse, Human, Mouse, Rabbit and Sheep Serum Proteins. |
| Form | Liquid (sterile filtered) |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target | Rat |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:15,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |