

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (Fc)-Alk. Phos., MinX Bo, Ho, Hu DNA-SEC-183864

Article Name	Goat F(ab)2 anti-Rat IgG (Fc)-Alk. Phos., MinX Bo, Ho, Hu
Biozol Catalog Number	DNA-SEC-183864
Supplier Catalog Number	SEC-183864
Alternative Catalog Number	DNA-SEC-183864
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA, IHC, WB
Species Reactivity	Rat
Immunogen	Rat IgG F(c) fragment
Conjugation	Alk. Phos.
Format	F(ab')2
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	Bovine, Equine, Human
Product Description	F(ab)2 Anti-Rat IgG F(c) Alkaline Phosphatase Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, considerat...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	F(ab')2 Anti-Rat IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Rat IgG, Rat IgG Fc and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat F(ab) or Bovine, Horse, and Human Serum Proteins.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl, 150 mM NaCl, 1 mM MgCl ₂ , 0.1 mM ZnCl ₂ , 50% (v/v) Glycerol, pH 8.0, sterile filtered, 0.05% NaN ₃
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	F(ab')2 Anti-Rat IgG Antibody has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code # NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for F(ab')2 Anti-Rat IgG Antibody.