

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 Anti-Rat IgG (H&L) Antibody Alkaline Phosphatase Conjugated (Min X Bv Hs Hu Ms Rb & Sh Serum Proteins) - 712-1533, AP, Polyclonal DNA-SEC-183866

Article Name	Goat F(ab)2 Anti-Rat IgG (H&L) Antibody Alkaline Phosphatase Conjugated (Min X Bv Hs Hu Ms Rb & Sh Serum Proteins) - 712-1533, AP, Polyclonal
Biozol Catalog Number	DNA-SEC-183866
Supplier Catalog Number	DNA-SEC-183866
Alternative Catalog Number	DNA-SEC-183866
Manufacturer	danova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rat
Immunogen	Rat IgG whole molecule
Conjugation	AP
Format	F(ab')2
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	Bovine,Equine,Human,Mouse,Rabbit,Sheep

Product Description	F(ab)2 Anti-Rat IgG Antibody is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays. F(ab)2 Anti-Rat IgG Antibody is ideal for investigators in Cancer, Immunology, and Micr...
Clonality	Polyclonal
Concentration	0.5 mg/mL
Isotype	Ig
Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Alkaline Phosphatase, anti-Goat Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) or Bovine, Horse, Human, Mouse, Rabbit and Sheep Serum Proteins.
Form	Liquid (sterile filtered)
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	F(ab)2 Anti-Rat IgG Antibody has been assayed against 1.0 ug of Rat IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for F(ab)2 Anti-Rat IgG Antibody.