

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Rat IgG (Fc)-Biotin, MinX Bo,Ho,Hu DNA-SEC-183867

Article Name	Goat F(ab)2 anti-Rat IgG (Fc)-Biotin, MinX Bo,Ho,Hu
Biozol Catalog Number	DNA-SEC-183867
Supplier Catalog Number	SEC-183867
Alternative Catalog Number	DNA-SEC-183867
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rat
Immunogen	Rat IgG F(c) fragment
Conjugation	Biotin
Format	F(ab')2
Target Specificity	IgG (Fc)
Cross-Adsorption (MinX)	Bovine,Equine,Human
Product Description	F(ab)2 Anti-Rat IgG F(c) Biotin Antibody was generated in goat and detects specifically Rat IgG F(c). Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be gi...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Rat IgG, Rat IgG F(c) and Rat Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Rat IgG F(ab) or Bovine, Horse and Human Serum Proteins.
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Rat
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. This product has been assayed against 1.0 µg of Rat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenzthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:15,000 to 1:90,000 of the reconstitution concentration is suggested for this product.