

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit Fab anti-Goat IgG (H+L)-Biotin, MinX none DNA-SEC-183926

| Article Name | Rabbit Fab anti-Goat IgG (H+L)-Biotin, MinX none |
|----------------------------|---|
| Biozol Catalog Number | DNA-SEC-183926 |
| Supplier Catalog Number | SEC-183926 |
| Alternative Catalog Number | DNA-SEC-183926 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Goat |
| Immunogen | Goat IgG whole molecule |
| Conjugation | Biotin |
| Format | Fab |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Fab Anti-Goat IgG Biotin Antibody generated in rabbit detects goat IgG. This product possesses the F(ab) region possessing the epitoperecognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are availa |
| Clonality | Polyclonal |

| Concentration | 0.5 mg/mL |
|--------------------|--|
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin and anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Rabbit IgG F(c). |
| Form | Lyophilized |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Goat |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |