

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Donkey Fab Anti-Goat IgG (H&L) Antibody Fluorescein Conjugated - 805-7202, FITC, Polyclonal DNA-SEC-183928**

Article Name	Donkey Fab Anti-Goat IgG (H&L) Antibody Fluorescein Conjugated - 805-7202, FITC, Polyclonal
Biozol Catalog Number	DNA-SEC-183928
Supplier Catalog Number	DNA-SEC-183928
Alternative Catalog Number	DNA-SEC-183928
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	WB
Species Reactivity	Goat
Immunogen	Goat IgG whole molecule
Conjugation	FITC
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Goat IgG Fluorescein Antibody generated in donkey detects goat IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are a...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Donkey Serum. No reaction was observed against anti-papain or anti-Donkey IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Goat
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 4.0, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	Fab Anti-Goat IgG Fluorescein Antibody has been tested by western blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.