

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Fab Anti-Guinea Pig IgG (H&L) Antibody Peroxidase Conjugated - 806-1302, HRP, Polyclonal DNA-SEC-183933

Article Name	Goat Fab Anti-Guinea Pig IgG (H&L) Antibody Peroxidase Conjugated - 806-1302, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183933
Supplier Catalog Number	DNA-SEC-183933
Alternative Catalog Number	DNA-SEC-183933
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Guinea pig
Immunogen	Guinea Pig IgG whole molecule
Conjugation	HRP
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Guinea Pig IgG Peroxidase Antibody generated in goat detects guinea pig IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibod...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Guinea Pig
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.