

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit Fab Anti-Human IgG (H&L) Antibody Peroxidase Conjugated - 809-4302, HRP, Polyclonal DNA-SEC-183954

Article Name	Rabbit Fab Anti-Human IgG (H&L) Antibody Peroxidase Conjugated - 809-4302, HRP, Polyclonal
Biozol Catalog Number	DNA-SEC-183954
Supplier Catalog Number	DNA-SEC-183954
Alternative Catalog Number	DNA-SEC-183954
Manufacturer	dianova
Host	Rabbit
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Human
Immunogen	Human IgG whole molecule
Conjugation	HRP
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Human IgG (H&L) Peroxidase Antibody generated in rabbit detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum im...

Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Rabbit IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Human
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000
Application Notes	Fab Anti-Human IgG (H&L) Peroxidase Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.