

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit Fab anti-Mouse IgG (H+L)-unconj., MinX none DNA-SEC-183960

| | |
|----------------------------|---|
| Article Name | Rabbit Fab anti-Mouse IgG (H+L)-unconj., MinX none |
| Biozol Catalog Number | DNA-SEC-183960 |
| Supplier Catalog Number | SEC-183960 |
| Alternative Catalog Number | DNA-SEC-183960 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA,IHC,WB |
| Species Reactivity | Mouse |
| Immunogen | Mouse IgG whole molecule |
| Conjugation | Unconjugated |
| Format | Fab |
| Target Specificity | IgG (H+L) |
| Cross-Adsorption (MinX) | no cross-adsorbtion |
| Product Description | Fab Anti-Mouse IgG (H&L) Antibody generated in rabbit detects Mouse IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are avail... |
| Clonality | Polyclonal |

| | |
|--------------------|---|
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Rabbit IgG F(c). |
| Form | Liquid (sterile filtered) |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target | Mouse |
| Antibody Type | Secondary Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |