

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat Fab anti-Rabbit IgG (H+L)-FITC, MinX none DNA-SEC-183965

Article Name	Goat Fab anti-Rabbit IgG (H+L)-FITC, MinX none
Biozol Catalog Number	DNA-SEC-183965
Supplier Catalog Number	SEC-183965
Alternative Catalog Number	DNA-SEC-183965
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	FLISA, IF
Species Reactivity	Rabbit
Immunogen	Rabbit IgG whole molecule
Conjugation	FITC
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Rabbit IgG (H&L) Antibody generated in goat detects rabbit IgG. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	FLISA Dilution: 1:10,000 - 1:50,000, Fluorochrome Protein Value: 2.0, IF Microscopy Dilution: 1:1,000 - 1:5,000
Application Notes	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.