

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Donkey Fab anti-Rabbit IgG (H+L)-HRPO, MinX none DNA-SEC-183970

Article Name	Donkey Fab anti-Rabbit IgG (H+L)-HRPO, MinX none
Biozol Catalog Number	DNA-SEC-183970
Supplier Catalog Number	SEC-183970
Alternative Catalog Number	DNA-SEC-183970
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Rabbit
Immunogen	Rabbit IgG whole molecule
Conjugation	HRPO
Format	Fab
Target Specificity	IgG (H+L)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Fab Anti-Rabbit IgG (H&L) antibody generated in Donkey detects specifically Rabbit IgG whole molecule. This Fab Anti-Rabbit IgG Secondary Antibody is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot,...
Clonality	Polyclonal

Concentration	1.0 mg/mL
Isotype	Ig
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Fab Anti-Rabbit IgG (H&L) was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Donkey Serum. No reaction was observed against anti-Papain or anti-Donkey IgG F(c).
Form	Lyophilized
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin
Target	Rabbit
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:8,000 - 1:32,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:3,000
Application Notes	Fab Anti-Rabbit IgG (H&L) Antibody is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.