

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD8a Antibody, Clone: [C8/468], Mouse, Monoclonal NSJ-V2043IHC-7ML

Artikelname	CD8a Antibody, Clone: [C8/468], Mouse, Monoclonal
Artikelnummer	NSJ-V2043IHC-7ML
Hersteller Artikelnummer	V2043IHC-7ML
Alternativnummer	NSJ-V2043IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human recombinant protein was used as the immunogen for this CD8a antibody.
Produktbeschreibung	CD8 is a cell surface receptor expressed either as a heterodimer with the beta chain or as a homodimer. A majority of thymocytes and a subpopulation of mature Tcells and NK cells express CD8a. It binds to MHC class 1 and through its association with ...
Klonalität	Monoclonal
Konzentration	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide, *For IHC use only*
Klon-Bezeichnung	[C8/468]
Isotyp	Mouse IgG1, kappa

NCBI	925
UniProt	P01732
Puffer	1X PBS, pH 7.4
Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Target-Kategorie	CD8A
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the CD8a antibody to be titrated up or down for optimal performance.>1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.