

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD6 Antibody, Clone: [C6/372], Mouse, Monoclonal NSJ-V2040-20UG

Article Name	CD6 Antibody, Clone: [C6/372], Mouse, Monoclonal
Biozol Catalog Number	NSJ-V2040-20UG
Supplier Catalog Number	V2040-20UG
Alternative Catalog Number	NSJ-V2040-20UG
Manufacturer	NSJ Bioreagents
Host	Mouse
Category	Antikörper
Application	IF, IHC-P
Species Reactivity	Human
Immunogen	Human recombinant CD6 protein was used as the immunogen for this antibody.
Product Description	CD6 is a type I transmembrane glycoprotein that contains a 24-amino acid signal sequence, three extracellular scavenger receptor cysteine-rich (SRCR) domains, a membrane-spanning domain and a 44-amino acid cytoplasmic domain. The CD6 glycoprotein is ...
Clonality	Monoclonal
Concentration	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Clone Designation	[C6/372]
Isotype	Mouse IgG1, kappa

NCBI	923
Buffer	1X PBS, pH 7.4
Purity	Protein G affinity chromatography
Form	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Target	CD6
Antibody Type	Primary Antibody
Application Dilute	Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Application Notes	The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the antibody to be titrated up or down for optimal performance. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.