

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rat Coagulation factor III/Tissue Factor protein, His tag, Unconjugated GTX00054-PRO

|                            |                                                                                                                                                                             |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name               | Rat Coagulation factor III/Tissue Factor protein, His tag, Unconjugated                                                                                                     |
| Biozol Catalog Number      | GTX00054-PRO                                                                                                                                                                |
| Supplier Catalog Number    | GTX00054-pro                                                                                                                                                                |
| Alternative Catalog Number | GTX00054-PRO-10                                                                                                                                                             |
| Manufacturer               | GeneTex                                                                                                                                                                     |
| Category                   | Proteine/Peptide                                                                                                                                                            |
| Application                | FA                                                                                                                                                                          |
| Species Reactivity         | Rat                                                                                                                                                                         |
| Conjugation                | Unconjugated                                                                                                                                                                |
| NCBI                       | <a href="#">25584</a>                                                                                                                                                       |
| UniProt                    | <a href="#">P42533</a>                                                                                                                                                      |
| Buffer                     | Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300. |
| Expression System          | E. coli                                                                                                                                                                     |
| Form                       | Lyophilized powder                                                                                                                                                          |
| Sequence                   | N-terminal His-Tag, Gly30~Glu252                                                                                                                                            |

Application Notes

TF (Tissue factor) also known for platelet tissue factor, factor III, thromboplastin, or CD142, is a cell surface glycoprotein present in subendothelial tissue and leukocytes. TF is necessary for the initiation of thrombin formation from the zymogen prothrombin. It has been proven that EGF can bind with the TF on the hemangioendotheliocytes. Thus a binding ELISA assay was conducted to detect the interaction of recombinant rat TF and recombinant rat EGF. Briefly, TF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl TF were then transferred to EGF-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-TF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of TF and EGF was in a dose dependent manner.